

P1 1140679

REC'D 08 MAR 2004

WIPO

PCT

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office

March 05, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/435,163

FILING DATE: December 20, 2002

RELATED PCT APPLICATION NUMBER: PCT/US03/41391

By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS



*L. Edelen*

L. EDELEN  
Certifying Officer

PRIORITY DOCUMENT  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

This Page Blank (uspto)

BEST AVAILABLE COPY

2/20/02  
JC904 U.S. PTO

12-23-02

Approved for use through 10/31/2002 OMB 0551-0032

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

# **PROVISIONAL APPLICATION FOR PATENT COVER SHEET**

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. **EV792050044**

JC906 U.S. PTO  
60/435163

INVENTOR(S)						
Given Name (first and middle (if any))		Family Name or Surname		Residence (City and either State or Foreign Country)		
Thomas J.		Smith		Weston, MA		
<input type="checkbox"/> Additional Inventors are being named on the _____ separately numbered sheets attached hereto						
TITLE OF THE INVENTION (500 characters max)						
LARGE CRYSTALLINE FORMULATIONS FOR SUSTAINED RELEASE DELIVERY OF ALKYLATING AGENTS						
Direct all correspondence to: CORRESPONDENCE ADDRESS						
<input type="checkbox"/> Customer Number _____ OR Type Customer Number here		<input type="checkbox"/> Firm or Individual Name Lisa M. Portscher PATENT TRADEMARK OFFICE				
Address		15 Braeburn Road				
Address						
City	Hyde Park	State	MA	ZIP	02136	
Country	USA	Telephone	617-361-5434	Fax		
ENCLOSED APPLICATION PARTS (check all that apply)						
<input checked="" type="checkbox"/> Specification Number of Pages		16		<input type="checkbox"/> CD(s), Number _____		
<input type="checkbox"/> Drawing(s) Number of Sheets				<input type="checkbox"/> Other (specify) _____		
<input type="checkbox"/> Application Data Sheet, See 37 CFR 1.76						
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT						
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.		FILING FEE AMOUNT (\$)				
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the filing fees						
<input type="checkbox"/> The Commissioner is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: _____		\$80.00				
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.						
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.						
<input checked="" type="checkbox"/> No.						
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____						

Respectfully submitted,

SIGNATURE Lisa M. Portscher

TYPED or PRINTED NAME Lisa M. Portscher

TELEPHONE 617-361-5434

Date 12/20/02

REGISTRATION NO.  
(if appropriate)  
Docket Number:

N/A

TJS-003

## **USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT**

This collection of information is required by 37 CFR 1.51. The information is used by the public to file (and by the PTO to process) a provisional application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete provisional application to the PTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, Washington, D.C. 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Box Provisional Application, Assistant Commissioner for Patents, Washington, D.C. 20231.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Approved for use through 10/31/2002. OMB 0651-0032  
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

# FEE TRANSMITTAL for FY 2003

Patent fees are subject to annual revision.

☒ Applicant claims small entity status. See 37 CFR 1.27

TOTAL AMOUNT OF PAYMENT (\$) 80.00

## Complete if Known

Application Number Unknown  
Filing Date 12/20/2002  
First Named Inventor Smith, Thomas J.  
Examiner Name Unknown  
Art Unit Unknown  
Attorney Docket No. TJS-003

## METHOD OF PAYMENT (check all that apply)

☒ Check ☐ Credit card ☐ Money Order ☐ Other ☐ None

☐ Deposit Account:

Deposit Account Number  
Deposit Account Name

The Commissioner is authorized to: (check all that apply)

☐ Charge fee(s) indicated below ☐ Credit any overpayments  
☐ Charge any additional fee(s) during the pendency of this application  
☐ Charge fee(s) indicated below, except for the filing fee to the above-identified deposit account.

## FEE CALCULATION

### 1. BASIC FILING FEE

Large Entity Fee Code (\$)	Small Entity Fee Code (\$)	Fee Description	Fee Paid
1001 740	2001 370	Utility filing fee	
1002 330	2002 165	Design filing fee	
1003 510	2003 255	Plant filing fee	
1004 740	2004 370	Reissue filing fee	
1005 160	2005 80	Provisional filing fee	80.00
SUBTOTAL (1) (\$)			80.00

### 2. EXTRA CLAIM FEES FOR UTILITY AND REISSUE

Total Claims  -20\*\* =  X  =   
Independent Claims  -3\*\* =  X  =   
Multiple Dependent  0 = 0

Large Entity		Small Entity		Fee Description
Fee Code	Fee (\$)	Fee Code	Fee (\$)	
1202	18	2202	9	Claims in excess of 20
1201	84	2201	42	Independent claims in excess of 3
1203	280	2203	140	Multiple dependent claim, if not paid
1204	84	2204	42	** Reissue independent claims over original patent
1205	18	2205	9	** Reissue claims in excess of 20 and over original patent

\*\*or number previously paid, if greater; For Reissues, see above

## FEE CALCULATION (continued)

### 3. ADDITIONAL FEES

Large Entity Fee Code (\$)	Small Entity Fee Code (\$)	Fee Description	Fee Paid
1051 130	2051 65	Surcharge - late filing fee or oath	
1052 50	2052 25	Surcharge - late provisional filing fee or cover sheet	
1053 130	1053 130	Non-English specification	
1812 2,520	1812 2,520	For filing a request for <i>ex parte</i> reexamination	
1804 920*	1804 920*	Requesting publication of SIR prior to Examiner action	
1805 1,840*	1805 1,840*	Requesting publication of SIR after Examiner action	
1251 110	2251 55	Extension for reply within first month	
1252 400	2252 200	Extension for reply within second month	
1253 920	2253 460	Extension for reply within third month	
1254 1,440	2254 720	Extension for reply within fourth month	
1255 1,960	2255 980	Extension for reply within fifth month	
1401 320	2401 160	Notice of Appeal	
1402 320	2402 160	Filing a brief in support of an appeal	
1403 280	2403 140	Request for oral hearing	
1451 1,510	1451 1,510	Petition to institute a public use proceeding	
1452 110	2452 55	Petition to revive - unavaldable	
1453 1,280	2453 640	Petition to revive - unintentional	
1501 1,280	2501 640	Utility issue fee (or reissue)	
1502 460	2502 230	Design issue fee	
1503 620	2503 310	Plant issue fee	
1460 130	1460 130	Petitions to the Commissioner	
1807 50	1807 50	Processing fee under 37 CFR 1.17(q)	
1806 180	1806 180	Submission of Information Disclosure Stmt	
8021 40	8021 40	Recording each patent assignment per property (times number of properties)	
1809 740	2809 370	Filing a submission after final rejection (37 CFR 1.129(a))	
1810 740	2810 370	For each additional invention to be examined (37 CFR 1.129(b))	
1801 740	2801 370	Request for Continued Examination (RCE)	
1802 900	1802 900	Request for expedited examination of a design application	

Other fee (specify)

\*Reduced by Basic Filing Fee Paid

SUBTOTAL (3) (\$) 0.00

## SUBMITTED BY

Name (Print/Type) Lisa M. Portscher  
Signature *Lisa M. Portscher*  
Registration No. (Attorney/Agent) N/A  
Telephone 617-361-5434  
Date 12/20/02

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

This collection of information is required by 37 CFR 1.17 and 1.27. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, Washington, DC 20231.

If you need assistance in completing the form, call 1-800-PTO-9199 (1-800-786-9199) and select option 2.

**LARGE CRYSTALLINE FORMULATIONS FOR  
SUSTAINED RELEASE DELIVERY OF ALKYLATING AGENTS**

**BACKGROUND OF THE INVENTION**

**Field of the Invention.**

This invention relates to the field of pharmaceutical sciences, and particularly to the formulation of sustained-release drug delivery systems for alkylating agents.

**Description of the Related Art.**

1. **Alkylating agents**

Cancer remains one of the leading causes of death worldwide. Current treatment of cancer utilizes surgery, radiation and drugs. Pharmaceutical treatment comprises agents which preferentially kill malignant cells. Among these drugs are the alkylating agents. Non-limiting examples of alkylating agents include the nitrogen mustards, alkyl sulfonate, nitrosurea, ethylenimine and methylmelamine, and triazene classes; cyclophosphamide, ifosfamide, thiotepa, melphalan, busulfan, carmustine, chlorambucil, hexamethylmelamine, and streptozocin; and the like and/or pharmaceutically acceptable salts thereof. Conditions treated with alkylating agents include: Hodgkin's disease and non-Hodgkin's lymphomas; acute and chronic lymphocytic leukemias; cancer of the lung, stomach, colon, pancreas, breast, ovary, cervix, bladder, and testis; multiple myeloma; Wilms' tumor; soft tissue sarcomas; primary macroglobulinemia; primary brain tumors; malignant melanoma; and malignant carcinoid

2. **Oral and Parenteral Formulations**

Alkylating agents are currently formulated for oral ingestion or parenteral injection. Therefore it is essential that the drugs dissolve into the gastro-intestinal or parenteral fluids in order to be absorbed into the body. For this reason, alkylating agents have been formulated as small particles that either have been mixed with binding agents and pressed into pills or tablets for oral administration, or have been dissolved in physiologically acceptable aqueous solvents for parenteral administration.

According to the Noyes Whitney Equation:

$$dc/dt = kS (C_s - C_t) \quad (1)$$

where  $dc/dt$  is the rate of dissolution,  $k$  is the dissolution rate constant,  $S$  is the surface area of the dissolving solid,  $C_s$  is the saturation concentration of drug in the diffusion layer and  $C_t$  is the concentration of drug in the dissolution media (or the bulk).

Of the variables in equation 1, only the surface area,  $S$ , is easily manipulated and, therefore, it is  $S$  which ultimately determines the rate of dissolution and drug delivery. In the prior art, alkylating agents have been formulated as small particles by milling in order to increase the surface area-to-volume ratio and therefore aid in rapid dissolution.

Furthermore, many drugs can be formulated as one of several polymorphs in order to alter their stability and dissolution properties. Polymorphism is the chemical characteristic of solid substances that can exist in one or more crystalline and/or amorphous forms. The different polymorphs are usually designated by Roman numerals, with the most stable polymorph under ambient conditions being designated as Form I. In order to increase their dissolution, some drugs are formulated as less stable polymorphs or as amorphous forms.

### 3. Sustained Release Formulations

In certain situations, it is desired that an alkylating agent be administered in a sustained release formulation, typically with the objectives of achieving a nearly constant, or zero order, release rate over a significant period of time and reducing the problem of non-compliance or non-adherence to therapy. Most sustained release systems employ a finely milled or micronized preparation of the active pharmaceutical ingredient as a starting point in the formulations. The release of the active pharmaceutical ingredient into the body is then controlled using matrices, membranes or other inactive ingredients or devices. Examples of methods and devices known in the art for sustained release formulations include liposomes, bioerodable matrices (e.g., PLA/PGLA matrices), drug-permeable implants (e.g., U.S. Pat. No. 3,993,073 to Zaffaroni), implants with drug-permeable and drug-impermeable membranes (e.g., U.S. Pat. No. 5,378,475 to

Smith et al.), and osmotic drug delivery systems (e.g., U.S. Pat. No. 4,439,196 to Higuchi).

The production and use of pharmaceutical preparations including larger drug particles to sustain delivery have been described in the art for certain antibiotics, insulin and steroids (see, e.g., Ansel, Allen and Popovich, eds., Pharmaceutical Dosage Forms and Drug Delivery Systems, 7<sup>th</sup> Edition, Lippincott Williams & Wilkins, Philadelphia, PA, 1999; Gennaro ed., Remington: The Science and Practice of Pharmacy, 19<sup>th</sup> Edition, Lippincott Williams & Wilkins: Philadelphia, PA., 1995), but the use of this technique has not been applied to the delivery of alkylating agents.

For example, suspensions of crystals of benzathine and procaine penicillin are useful in the treatment of rheumatic fever and other infections. Typically, therapeutic levels of these antibiotics can be sustained for 14-28 days after a single IM injection (Cadorniga et al. (1991), *Eur. J. Drug Metab. Pharmacokinet.* 3:379-84; Kaplan et al. (1989), *J. Pediatr.* 115(1):146-50; U.S. Pat. No. 2,627,491; U.S. Pat. No. 2,515,898).

Similarly, suspensions of crystals of zinc insulins have slow release properties that depend on crystal size (Nichols (2000), in Remington: The Science and Practice of Pharmacy, Gennaro, ed., Lippincott Williams & Wilkins: Philadelphia, PA. p. 1371). The principal determinant of sustained antiglycemic action in these slow-release or "lente" insulin preparations is the dissolution of the insulin crystals.

Sustained release suspensions of corticosteroids for local delivery (intralesional, intra-articular) or systemic delivery achieve their duration of action due to slow dissolution of crystals of insoluble salts (Mollmann et al. (1977), *Fortschr. Med.* 95(14): 972-8).

There remains a need, however, for new and improved methods and products for the sustained release delivery of alkylating agents.

#### SUMMARY OF THE INVENTION

The present invention is generally directed to the production and use of pharmaceutical preparations including large and/or shape-controlled coated or uncoated crystalline formulations for the purpose of sustained-release delivery of alkylating agents. The invention depends, in part, upon the recognition that, if a larger crystal of a drug is

introduced, there will be a sustained release effect due to the decreased surface area-to-volume ratio of the larger particles. In addition, if the particle is formulated to be substantially flat, then the kinetics of drug release can be made to approach zero order or linear delivery. Moreover, formulation using the most stable polymorph (Form I) will aid in sustaining release because of decreased solubility from the surface, and will also increase the chemical stability of the active pharmaceutical ingredient. Furthermore, the coating of the particle with a semi-permeable membrane can cause the delivery rate to be predicated by the difference in the concentration of drug in solution between the inside and outside of the coating. The chemical potential difference between the inside and outside will be constant as long as the solution is saturated inside and at a "sink" concentration outside. In the present invention, such conditions will apply for the majority of the drug delivery time period, therefore causing sustained pseudo-zero order delivery.

In some embodiments of the invention, the crystals have at least one linear dimension greater than 50 $\mu$ m, greater than 100 $\mu$ m, greater than 500 $\mu$ m, or greater than 1mm. In some embodiments in which the formulations are heterogeneous with respect to crystal size, at least 70%, 80%, 90% or 95% of the crystals have at least one linear dimension greater than 50 $\mu$ m, greater than 100 $\mu$ m, greater than 500 $\mu$ m, or greater than 1mm.

In some embodiments, the formulations are produced as bodies which are relatively thin in one dimension relative to perpendicular dimensions, such as substantially flat or disc-shaped bodies.

In some embodiments, the active pharmaceutical ingredient is crystallized in a stable crystalline form, or the most stable of multiple crystalline forms.

In some embodiments, the pharmaceutical preparation of the invention is a surgically implantable or injectable (via a needle/trochar) implant.

In some embodiments, the pharmaceutical preparation of the invention is an injectable suspension.

In some embodiments, the crystals are uncoated.

In some embodiments, the crystals are coated with a biodegradable semi-permeable polymer coating. In specific embodiments, the coating can be PLA; PLGA; a polyorthoester; or a polyanhydride.

In some embodiments, the crystals are coated with a biocompatible semi-permeable polymer coating. In specific embodiments, the coating can be PVA; EVA; or silicone.

In another aspect, the invention provides methods for treating a mammal suffering from or at risk of a condition for which administration of an alkylating agent is indicated. The methods involve administering a pharmaceutical preparation of the invention to the mammal such that an effective amount of the active pharmaceutical ingredient is provided over a sustained period of time.

In another aspect, the invention provides a pharmaceutical preparation and kits comprising large crystalline particles. In some embodiments, the large crystalline particles are in a dry form that can be suspended in a separately provided diluent prior to use. In some embodiments, a kit is provided containing separate containers of the large crystalline particles in a dry form and a diluent for forming the suspension prior to use. In other embodiments, a multi-chambered product is provided in which the large crystalline particles are contained in one chamber in a dry form and a diluent is contained in another chamber, such that a suspension is formed when the wall separating the chambers is ruptured and the particles and diluent are mixed.

### DETAILED DESCRIPTION

#### General Considerations.

The present invention is generally directed to the production and use of pharmaceutical preparations including large crystalline formulations for the purpose of sustained-release delivery of alkylating agents. In contrast to the teachings of the prior art, the present invention is dependent, in part, upon the discovery that large crystalline formulations of these active pharmaceutical ingredients, employing crystals having linear dimensions exceeding 50 $\mu$ m, 100 $\mu$ m, 500 $\mu$ m, or even 1mm, are useful for sustained-release drug delivery. Moreover, it has been discovered that such large crystalline



formulations can be produced in flattened shapes (i.e., shapes in which the body is substantially smaller in one linear dimension relative to perpendicular dimensions) which dissolve with pseudo-zero order kinetics. Furthermore, the coating of such crystals can enhance the linearity of release. Some active pharmaceutical ingredients can exist in multiple crystalline forms. For such active ingredients, it has been discovered that the stable or most stable form is the most useful for sustained release drug delivery.

#### Definitions.

All technical and scientific terms used herein, unless otherwise defined below, are intended to have the same meaning as commonly understood by one of ordinary skill in the art; references to techniques employed herein are intended to refer to the techniques as commonly understood in the art, including variations on those techniques or substitutions of equivalent techniques which would be apparent to one of skill in the art. In order to more clearly and concisely describe the subject matter which is the invention, the following definitions are provided for certain terms which are used in the specification.

The term "pharmaceutically acceptable carrier" as used herein means a material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting the active pharmaceutical ingredients from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "pharmaceutically acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient (i.e., suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problems or complications, commensurate with a reasonable benefit/risk ratio).

As used herein, the term "effective amount" of an active pharmaceutical ingredient means the total amount of the active pharmaceutical ingredient in a composition that is sufficient to cause a statistically significant change of a detectable biochemical or phenotypic characteristic. When applied to an individual active pharmaceutical ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active

pharmaceutical ingredients that result in the effect, whether administered in combination, serially or simultaneously.

As used herein, the term "substantially pure" means a preparation which contains at least 60% (by dry weight) of the active pharmaceutical ingredient of interest, exclusive of the weight of other intentionally included compounds. In certain embodiments, the preparation is at least 75%, at least 90%, or at least 99% the active pharmaceutical ingredient of interest by dry weight, exclusive of the weight of other intentionally included compounds. Purity can be measured by any appropriate method, e.g., column chromatography, gel electrophoresis, amino acid compositional analysis or HPLC analysis. If a preparation intentionally includes two or more different active pharmaceutical ingredients of the invention, a "substantially pure" preparation means a preparation in which the total dry weight of the active pharmaceutical ingredients of the invention is at least 60% of the total dry weight, exclusive of the weight of other intentionally included compounds. For preparations containing two or more active pharmaceutical ingredients of the invention, the total weight of the active pharmaceutical ingredients of the invention should be at least 75%, at least 90%, or at least 99%, of the total dry weight of the preparation, exclusive of the weight of other intentionally included compounds. Thus, if the active pharmaceutical ingredients of the invention are mixed with one or more other compounds (e.g., diluents, stabilizers, detergents, excipients, salts, sugars, lipids) for purposes of administration, stability, storage, and the like, the weight of such other compounds is ignored in the calculation of the purity of the preparation.

As used herein, the term "mammalian subject" means any member of the class Mammalia including, without limitation, humans and nonhuman primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs, and the like. The term does not denote a particular age or sex. Thus, adult and newborn subjects, as well as fetuses, whether male or female, are intended to be covered.

As used herein, the terms "modulate" or "affect" mean to either increase or decrease. As used herein, the terms "increase" and "decrease" mean, respectively,

statistically significantly increase (*i.e.*,  $p < 0.5$ ) and statistically significantly decrease (*i.e.*,  $p < 0.5$ ).

As used herein, the recitation of a numerical range for a variable is intended to convey that the invention may be practiced with the variable equal to any of the values within that range. Thus, for a variable which is inherently discrete, the variable can equal each integer value of the numerical range, including the end-points of the range. Similarly, for a variable which is inherently continuous, the variable can equal each real value of the numerical range, including the end-points of the range. As an example, a variable which is described as having values between 0 and 2, can be 0, 1 or 2 for variables which are inherently discrete, and can be 0.0, 0.1, 0.01, 0.001, or any other real value  $\leq 2$  for variables which are inherently continuous.

As used herein, unless specifically indicated otherwise, the word "or" is used in the "inclusive" sense of "and/or" and not the "exclusive" sense of "either/or."

#### Crystal Size.

In one aspect, the invention provides pharmaceutical formulations in which the active pharmaceutical ingredient is in the form of large crystals. This distinguishes the formulations of the invention from prior art formulations in which active ingredients are machined or milled to produce small or micronized crystals of the drug, which then must be combined with matrices, semi-permeable membranes, pumps or other inactive ingredients or devices in order to achieve the effect of sustained release delivery.

In some embodiments, the crystals have at least one linear dimension greater than 50 $\mu$ m, greater than 100 $\mu$ m, greater than 500 $\mu$ m, or greater than 1mm. In some embodiments in which the formulations are heterogeneous with respect to crystal size, at least 70%, 80, 90% or 95% of the crystals have at least one linear dimension greater than 50 $\mu$ m, greater than 100 $\mu$ m, greater than 500 $\mu$ m, or greater than 1mm.

#### Body Shape.

The shape of a solid formulation, including implants or solid bodies that may be present in a liquid suspension, can be described by reference to three perpendicular or orthogonal axes. In some embodiments, the formulations are produced as bodies which

are substantially smaller in one dimension relative to perpendicular dimensions, such as substantially flat or disc-shaped bodies. Such bodies will present a more constant surface area over time during dissolution. Because dissolution occurs at the surface of the body, such relatively constant surface areas are useful to approximate or achieve pseudo-zero order kinetics of delivery.

#### Crystal Form.

Polymorphism is the chemical characteristic of solid substances that can exist in one or more crystalline and/or amorphous forms. The different polymorphs are usually designated by Roman numerals, with the most stable polymorph under ambient conditions being designated as Form I.

In some embodiments, the active pharmaceutical ingredient is crystallized in a stable crystalline form, or the most stable of multiple crystalline forms (e.g., Form I). Because this crystalline form tends to be the least soluble in water, it has been avoided in the prior art. However, it has been discovered that utilizing the most stable polymorph can significantly enhance the ability to achieve sustained release rates.

#### Active Pharmaceutical Ingredients.

Active pharmaceutical ingredients which may be formulated according to the invention include any alkylating agent drug which may be crystallized according to the invention. Non-limiting examples of alkylating agents include the nitrogen mustards, alkyl sulfonate, nitrosurea, ethylenimine and methylmelamine, and triazene classes; cyclophosphamide, ifosamide, thiotepa, melphalan, busulfan, carmustine, clorambucil, hexamethylmelamine, and streptozocin; and the like and/or pharmaceutically acceptable salts thereof.

#### Production of Large Crystalline Formulations.

The large crystalline formulations of the invention, although not produced in the prior art, can be produced by methods well known in the art. The active pharmaceutical

ingredients in the formulations can be any alkylating agent for which the large crystalline formulations of the invention can be produced.

#### 1. Solvation Crystallization.

A substantially pure pharmaceutical preparation of the alkylating agent drug is synthesized, isolated or otherwise obtained. The purity is tested to ensure suitability for use in a pharmaceutical preparation.

The active pharmaceutical ingredient is dissolved in an appropriate volatile solvent to form a solution. The solution is then dispensed into properly shaped drying areas, such as flat-bottomed wells or flat or dimpled surfaces. The solvent is then allowed to evaporate at an appropriate rate (e.g., controlling the rate by controlling the temperature and/or the humidity and/or the partial pressure of the solvent or other gases in the ambient atmosphere) to obtain crystals of the desired size.

Alternatively the solution of active pharmaceutical ingredient in solvent is sprayed onto a cooling slab surface. A disc is formed by evaporation of the solvent and the disc parameters (e.g., size, shape, weight) are adjusted by adjusting the parameters of the spraying step (e.g., volume of drops, concentration, ambient pressure and temperature, slab temperature, velocity of spray, nozzle size, etc.).

The resultant crystals are sorted by size and/or subjected to mechanical forces to modify the size or shape of the crystalline bodies. For example, the bodies can be passed through sieves to isolate bodies within desired size ranges, and can be ground, lathed, milled, cut (e.g., with a laser), etc., to reduce the size of alter the shape of the bodies.

#### 2. Melting and Recrystallization.

A substantially pure pharmaceutical preparation of the alkylating agent drug is synthesized, isolated or otherwise obtained. The purity is tested to ensure suitability for use in a pharmaceutical preparation.

The melting point of the active pharmaceutical ingredient is determined by standard methods (e.g., visual inspection while slowly increasing or decreasing ambient temperature and/or by reference to the literature). For each active pharmaceutical ingredient, the optimum melting process is that which melts the drug completely while

minimizing molecular breakdown. This optimum process can be determined for each active pharmaceutical ingredient by one skilled in the art through the process of varying key parameters in the melting process (e.g., rate, intensity, agitation or mixing, etc.). The results of the process can be assessed by determining the purity of the resulting melted form (i.e., lack of molecular breakdown) through HPLC or NMR analysis. After melting, the active pharmaceutical ingredient is cooled, actively or passively, so as to promote the formation of large crystals of the desired crystalline form (e.g., Form I). Optimum cooling parameters and process (e.g., mixing, rate of cooling, final temperature, cooling substrate or surface characteristics, etc.) can be determined by varying these parameters experimentally and characterizing the resultant crystals in terms of size, shape and polymorph state.

The resultant crystals can then be sorted by size and/or subjected to mechanical forces to modify the size or shape of the crystalline bodies. For example, the bodies can be passed through sieves to isolate bodies within desired size ranges, and can be ground, lathed, milled, cut (e.g., with a laser), etc., to reduce the size or alter the shape of the bodies.

### 3. Seed Crystallization.

A substantially pure pharmaceutical preparation of the alkylating agent is synthesized, isolated or otherwise obtained. The purity is tested to ensure suitability for use in a pharmaceutical preparation.

The active pharmaceutical ingredient is dissolved in an appropriate solvent to achieve a saturated solution. The saturated solution is then seeded with a crystal of the active pharmaceutical ingredient to initiate crystal formation from the saturated solution. The crystals are then harvested (e.g., by straining from solution and drying, or by evaporating the solvent).

The resultant crystals can then be sorted by size and/or subjected to mechanical forces to modify the size or shape of the crystalline bodies. For example, the bodies can be passed through sieves to isolate bodies within desired size ranges, and can be ground, lathed, milled, cut (e.g., with a laser), etc., to reduce the size of alter the shape of the bodies.

#### 4. Pressure.

A substantially pure pharmaceutical preparation of the alkylating agent is synthesized, isolated or otherwise obtained. The purity is tested to ensure suitability for use in a pharmaceutical preparation.

The active pharmaceutical ingredient in powder form is compressed under high pressure (e.g., for 10 to 100 M tons per  $\text{cm}^2$ ; 25 to 50 M tons per  $\text{cm}^2$ ) for a time sufficient for a glass transformation to take place.

The resultant discs can then be sorted by size and/or subjected to mechanical forces to modify the size or shape of the crystalline bodies. For example, the bodies can be passed through sieves to isolate bodies within desired size ranges, and can be ground, lathed, milled, cut (e.g., with a laser), etc., to reduce the size of alter the shape of the bodies.

#### Coatings.

Standard coating techniques well known in the art can be used to coat the large crystals. Examples include dipping, pan coating and spray coating. Care must be taken that the active pharmaceutical ingredient is not soluble in the solvent used in the polymeric coating. As discussed below, the coating can be a biodegradable polymer, a biocompatible polymer, or a biodegradable biocompatible polymer.

#### Biodegradable Polymers.

In some of the implementations of this invention, the large crystalline particles or implants are coated with biodegradable polymers. These polymers are well known in the art and examples include: naturally occurring polymers such as sugar phosphates, which are known to be biodegradable, and synthetic polymers such as polylactides and polyglycolic acids, which are also biodegradable. Lactic acid copolymers offer a degree of flexibility in choosing the life of a polymer matrix, because the half-life can be controlled by varying the amount and type of co-monomer used. Examples of suitable copolymers are glycolide,  $\beta$ -propiolactone, tetramethylglycolide,  $\beta$ -butyrolactone, tetramethylglycolide,  $\beta$ -butyrolactone,  $\gamma$ -butyrolactone, pivalolactone, intramolecular

cyclic esters of  $\alpha$ -hydroxybuteric acid,  $\alpha$ -hydroxy, isovaleric acid,  $\alpha$ -hydroxycaproic acid,  $\alpha$ -hydroxy ethylbuteric acid,  $\alpha$ -hydroxy isocaproic,  $\alpha$ -hydroxy  $\beta$ -methyl valeric acid,  $\alpha$ -hydroxy heptonic acid,  $\alpha$ -hydroxy octanic acid,  $\alpha$ -hydroxy deccanoic acid,  $\alpha$ -hydroxy myristic acid,  $\alpha$ -hydroxy stearic acid,  $\alpha$ -hydroxy ligocenic acid, and  $\beta$ -phenol lactic acid. As also known in the art, polyglycolic acids can provide excellent biodegradable properties.

#### Biocompatible Polymers.

In some of the implementations of this invention, the large crystalline particles or implants are coated with biocompatible polymers. These polymers are well known in the art and examples include: acyl substituted cellulose acetates and alkyl derivatives thereof, partially and completely hydrolyzed alkylene-vinyl acetate copolymers; unplasticized polyvinyl chloride; cross-linked homo- and copolymers of polyvinyl acetate; cross-linked polyesters of acrylic and methacrylate; polyvinyl alkyl ethers; polyvinyl fluoride; silicone; polycarbonate; polyurethane; polyamide; polysulphones; styrene acrylonitrile copolymers; cross-linked poly(ethylene oxide); poly(alkylenes); poly(vinyl imidazole); poly(esters); poly(ethylene terephthalate); and chlorosulphonated polyolefins.

#### Dosage Forms.

In another aspect, the invention provides pharmaceutical preparations comprising one or more of the alkylating agents described above, formulated together with one or more pharmaceutically acceptable carriers (additives) and/or diluents. As described in detail herein, the pharmaceutical compositions of the invention are specially formulated for administration in the form of large crystalline particles in suspension or in the form of implants. These formulations are administered by parenteral administration, for example, by subcutaneous or intramuscular injection for suspensions, or by subcutaneous or intramuscular implantation for implants. Implants or suspensions may be designed for local, or targeted, delivery or for systemic delivery of the drug.

Because of the potential for embolic sequelae, intravenous or intra-arterial injections may not be appropriate, and subcutaneous (SC) or intramuscular (IM)



injections may be safer. An injection volume of 0.1 to 10 cc, or 1 to 6 cc, can be used for IM injection, and a volume of 0.1 to 2 cc can be used for SC.

In some embodiments of the invention, the crystals of the alkylating agent drug are suspended in a pharmaceutically acceptable carrier which serves as a dispersion medium to form a suspension for injection into a subcutaneous or intramuscular position for sustained pseudo-linear delivery over a period of greater than 1 week, greater than 2 weeks, greater than 4 weeks, or greater than 12 weeks.

Suspensions of large particles will tend to settle more rapidly than suspensions of small particles. Therefore, in some embodiments, a commercial product of the invention can be distributed as large crystalline particles in a dry form that can be suspended in a separately provided diluent prior to use. In other embodiments, a kit is provided containing separate containers of the large crystalline particles in a dry form and a diluent for forming the suspension prior to use. Alternatively, a multi-chambered product can be provided in which the large crystalline particles are contained in one chamber in a dry form and a diluent is contained in another chamber, such that a suspension is formed when the wall separating the chambers is ruptured and the particles and diluent are mixed. In addition to the diluent, appropriate suspending agents and surfactants well known in the art can be included.

When administered in the form of a suspension, a liquid carrier such as water, physiological saline solution, petroleum, oil (e.g., animal, plant or mineral oils, such as cottonseed, groundnut, corn, germ, olive, castor, peanut, mineral, soybean, sesame, or synthetic oils) can be used. The suspension form of the pharmaceutical can further contain inert diluents commonly used in the art, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, ethyl oleate, benzyl alcohol, benzyl benzoate, glycols (e.g., ethylene glycol, propylene glycol, polyethylene glycol, 1,3-butylene glycol), glycerol, tetrahydrofuryl alcohol, dextrose or other saccharide solutions, and mixtures thereof. Suspensions, in addition to the active compounds, can contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan fatty acid esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof. When administered in suspension form, the

pharmaceutical composition can contain about 0.01 to 99% by weight of the crystals, 0.1 to 90% of the crystals, or about 1 to 50% of the crystal.

Pharmaceutically acceptable carriers can include solutions or mixtures of compounds such as sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; gelatin; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; alginic acid; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; and other non-toxic compatible substances routinely employed in pharmaceutical formulations.

In some embodiments of the invention, the crystals of the alkylating agent drug are molded, shaped, compressed or formed into a solid implant for implantation into a subcutaneous or intramuscular position for sustained pseudo linear delivery over a period of greater than 1 week, greater than 2 weeks, greater than 4 weeks, or greater than 12 weeks. Such implants can include (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, hydroxypropylmethyl cellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; or (3) embedding compositions, such as polymeric substances and waxes.

#### Methods of Treatment

In another aspect, the invention provides methods for treating a mammal suffering from or at risk of a condition for which administration of an alkylating agent is indicated. Such conditions include: Hodgkin's disease and non-Hodgkin's lymphomas; acute and chronic lymphocytic leukemias; cancer of the lung, stomach, colon, pancreas, breast, ovary, cervix, bladder, and testis; multiple myeloma; Wilms' tumor; soft tissue sarcomas; primary macroglobulinemia; primary brain tumors; malignant melanoma; and malignant carcinoid.

In general, the methods involve administering a pharmaceutical preparation of the invention to the mammal such that an effective amount of the active pharmaceutical

ingredient is provided over a sustained period of time. The mammals can be humans, or can be nonhuman primates such as chimpanzees or other apes or monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs, and the like. The delivery may be systemic or targeted to a specific location or lesion.

The sustained release formulation can be a coated or uncoated large crystal, or a coated or uncoated implant. In those embodiments in which the crystals or implants are coated, the coating can be a biodegradable polymer and/or a biocompatible polymer. In those embodiments in which the formulation includes large crystals, the crystals can be held in a suspension for injection.

In some embodiments, the formulations will be designed to provide sustained pseudo-linear delivery over a period of greater than 1 week, greater than 2 weeks, greater than 4 weeks, or greater than 12 weeks. The amount of the drug and the rate of delivery can be determined by the physician and/or pharmacist and will depend upon such factors as the potency of the active pharmaceutical ingredient, the severity of the condition, and the age, sex and weight of the subject. The rate of drug delivery is determined to provide an effective amount of the active pharmaceutical ingredient over a sustained period of time.

#### EQUIVALENTS

While this invention has been particularly shown and described with references to certain embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the invention. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described specifically herein. Such equivalents are intended to be encompassed in the scope of the invention.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**